

DecaLabel™ DNA Labeling Kit (#K0622, for 30 reactions)

Labeling of DNA by random oligonucleotide-primed synthesis is based on the investigations of A. Feinberg and B. Vogelstein [1, 2] and is a good alternative to nick translation for producing uniformly radioactive DNA of high specific activity. The method relies on priming of the polymerase reaction on the template DNA with random decanucleotide primers. The complementary strand is synthesized from the 3'-end of the primer with the help of the Large Fragment of DNA Polymerase I, Exonuclease Minus (Klenow Fragment, exo^-) in the presence of labeled deoxyribonucleoside triphosphates. Using Klenow fragment, exo^- the reaction time can be prolonged without the fear of labeled probe degradation. The DecaLabel™ DNA Labeling Kit offered by Fermentas is an efficient and convenient means for preparation of highly labeled DNA ($>10^9$ dpm/ μ g) for use diverse procedures in molecular biology, such as various types of hybridization analyzes [3-6]. Decanucleotides are more efficient primers than hexanucleotides, and the optimal reaction time is 5min. Either dATP or dCTP could be used with the kit as a radioactive precursor.

COMPONENTS OF THE KIT

1. Klenow Fragment, exo^- , 5u*/ μ l:

30 μ l of the enzyme solution in buffer containing 50% glycerol.

2. Decanucleotide in 5x Reaction Buffer:

300 μ l of 0.25M Tris-HCl buffer (pH 8.0 at 20°C) containing 25mM MgCl₂, 5mM dithiothreitol and Random (decamer) primer (12.5o.u./ml).

3. Mix A (minus dATP):

90 μ l of 0.33mM dGTP, 0.33mM dTTP, 0.33mM dCTP aqueous solution.

4. Mix C (minus dCTP):

90 μ l of 0.33mM dGTP, 0.33mM dATP, 0.33mM dTTP aqueous solution.

5. dNTP Mix:

120 μ l of 0.25mM dGTP, 0.25mM dATP, 0.25mM dTTP, 0.25mM dCTP aqueous solution.

6. Control Template:

for 5 control labeling reactions, 20 μ l of λ phage DNA/HindIII fragments (10 μ g/ml).

7. Water, nuclease-free:

1.5ml of 0.22 μ m membrane-filtered molecular biology grade water.

One unit of Klenow fragment, exo^- catalyzes the incorporation of 10 nmoles of deoxyribonucleotides into a polynucleotide fraction (adsorbed on DE-81) in 30min at 37°C, using poly(dA-dT)•poly(dA-dT) as a template•primer.

Radioactive DNA Labeling

- ➊ Add the following components into 1.5ml microcentrifuge tube:

| | |
|--------------------------------------|----------|
| DNA template (100ng) | 10μl, |
| Decanucleotide in 5x Reaction Buffer | 10μl, |
| Water, nuclease-free | to 40μl. |

 Vortex the tube and spin down in a microcentrifuge for 3-5sec. Incubate the tube in a boiling water bath for 5-10min and cool it on ice. Spin down quickly.
- ➋ Based on your choice of labeled triphosphate (dATP or dCTP) use Mix A or Mix C, respectively.
- ➌ Add the following components in the same tube:

| | |
|--|------|
| Mix A (or Mix C) | 3μl, |
| [α- ³² P]-dATP (or [α- ³² P]-dCTP)** (1.85MBq = 50μCi) | 6μl, |
| Klenow fragment, exo ⁻ (5u) | 1μl. |

 Shake the tube and spin down in a microcentrifuge for 3-5sec. Incubate for 5min at 37°C.
- ➍ Add 4μl of dNTP Mix and incubate for 5min at 37°C.
- ➎ Stop the reaction by the addition of 1μl 0.5M EDTA, pH 8.0.
- ➏ The labeled DNA is used directly for hybridization or stored at -20°C. Removal of the unincorporated label is not necessary for most applications, since the levels of its utilization are usually high. If required, the unincorporated dNTP can be removed by chromatography on Sephadex[®] G-50 or by selective precipitation of DNA with ethanol in the presence of ammonium acetate [7].

** If you use ethanol solution of labeled [α-³²P]-dATP or [α-³²P]-dCTP, the needed quantity of these nucleotides dry under vacuum and redissolve in 6μl of Water, nuclease-free.

Determination of the Percentage of Label Incorporation

The percentage of incorporation is determined by DE-81 filter-binding assay.

- ➊ Dilute 1μl of the reaction mix 1:100 with 0.2M EDTA or water. Spot 5μl of it on each of the two Whatman[®] DE-81 filters (1.5 x 1.5cm).
- ➋ Dry the filters under a heat lamp. Keep one filter aside and use it directly for the determination of total dpm in the sample.
- ➌ Wash the other filter 3 times for 5min in 10ml 7.5% (w/v) Na₂HPO₄ · 12H₂O for the removal of the unincorporated dNTPs, then once in water and in acetone.
- ➍ Dry the washed filter under a heat lamp.
- ➎ Transfer and count the filters in an appropriate radioactivity counter.
- ➏ The percentage of label incorporation into DNA is defined as below:

$$\frac{\text{incorporated radioactivity (washed filter)}}{\text{total radioactivity (unwashed filter)}} \times 100\%$$

- ➐ Calculate the specific activity of the probe: for the calculation of the specific activity of the labeled DNA probe, it is first necessary to evaluate the theoretical amount of DNA generated in the reaction assuming that 100% incorporation of radioactivity occurred:

$$\frac{\mu\text{Ci dNTP added} \times 4 \times 330\text{ng/nmole}}{\text{specific activity dNTP (Ci/mmmole} = \mu\text{Ci/nmole)}} = \text{ng theoret. yield}$$

330ng/nmole = average molecular weight of a nucleotide.

Further, calculate the percentage incorporation according to DE-81 filter-binding assay results:

$$\frac{\text{dpm incorporated}}{\text{total dpm}} \times 100\% = \% \text{ incorporation}$$

Determine the amount of DNA synthesized:

$$\% \text{ incorporation} \times 0.01 \times \text{theoret. yield} = \text{ng DNA synthesized}$$

Calculate the specific activity of the product:

$$\frac{\text{total dpm incorp.} (\text{dpm incorp.} \times 20 \times 50)}{(\text{ng DNA synth.} + \text{ng input DNA}) \times 0.001 \mu\text{g/ng}} = \text{dpm}/\mu\text{g}$$

Note. The factors 20 and 50 are derived while using 5 μ l of 1:100 dilution for DE-81 filter-binding assay and converting this back to 50 μ l total reaction volume.

Example. If you use 50 μ Ci (1.85 MBq) [α -³²P]-dCTP (3000Ci/mmol) in a standard reaction, the calculation is as follows:

$$\frac{50 \mu\text{Ci} \times 4 \times 330 \text{ ng/nmole}}{3000 \mu\text{Ci/nmole}} = 22 \text{ ng theoretical yield}$$

Assuming that 14.57 x 10⁴dpm remained on the washed DE-81 filter and the unwashed filter retained 17.32 x 10⁴dpm, we recur:

$$\frac{14.57 \times 10^4}{17.32 \times 10^4} \times 100\% = 84.1\% \text{ incorporation}$$

$$0.841 \times 22 \text{ ng} = 18.5 \text{ ng DNA synthesized}$$

Thus, the specific activity is:

$$\frac{14.57 \times 10^4 \text{ dpm} \times 20 \times 50}{(18.5 \text{ ng} + 100 \text{ ng}) \times 0.001 \mu\text{g/ng}} = 1.23 \times 10^9 \text{ dpm}/\mu\text{g}$$

QUALITY CONTROL

All components of the kit are tested in a labeling reaction of λ DNA/HindIII fragments, obtained specific activity 1.2x10⁹ dpm/ μ g DNA.

Quality authorized by:



Laima Kazilioniene

References

1. Feinberg, A.P., Vogelstein, B., Biochem. 132, 6-13, 1983.
2. Feinberg, A.P., Vogelstein, B., Biochem. 137, 266-267, 1984.
3. Haase, A. et al., Virology 7, 189-226, 1984.
4. Meinkoth, M., Wahl, G., Anal. Biochem. 158, 267-284, 1984.
5. Grunstein, M., Hogness, D.S., Proc. Natl. Acad. Sci. USA 72, 3961-3965, 1975.
6. Southern, E.M., J. Mol. Biol., 98, 503-517, 1975.
7. Sambrook, J., Fritsch, E.F. and Maniatis, T., Molecular Cloning: A Laboratory Manual; Second Edition, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y., 1989.

Trademarks

DecaLabel is a trademark of Fermentas.

Sephadex is a registered trademark of Pharmacia Biotech, Inc.

Whatman is a registered trademark of Whatman Ltd.

PRODUCT USE LIMITATION.

This product is developed, designed and sold exclusively for research purposes and *in vitro* use only. The product was not tested for use in diagnostics or for drug development, nor is it suitable for administration to humans or animals. Please refer to www.fermentas.com