



CERTIFICATE OF ANALYSIS

Eco52I (EagI)

#ER0332 2500 u

Lot: Expiry Date:

5'...C↓G G C C G...3'
3'...G C C G G↑C...5'

Concentration: 10 u/μl
Source: *Escherichia coli* RFL52
Supplied with: 1 ml of 10X Buffer Eco52I
 1 ml of 10X Buffer Tango™

Store at -20°C



In total 3 vials.

BSA included: Lot# BSA62-313P

RECOMMENDATIONS

1X Buffer Eco52I (for 100% Eco52I digestion)

10 mM Tris-HCl (pH 8.5), 3 mM MgCl₂, 100 mM NaCl,
0.1 mg/ml BSA.

Incubation temperature

37°C.

Unit Definition

One unit is defined as the amount of Eco52I required to digest 1 μg of lambda DNA-Eco81I fragments in 1 hour at 37°C in 50 μl of recommended reaction buffer.

Dilution

Dilute with Dilution Buffer (#B19): 10 mM Tris-HCl (pH 7.4 at 25°C), 100 mM KCl, 1 mM EDTA, 1 mM DTT, 0.2 mg/ml BSA and 50% glycerol.

Double Digests

Tango™ Buffer is provided to simplify buffer selection for double digests. 98% of Fermentas restriction enzymes are active in a 1X or 2X concentration of Tango™ Buffer. Please refer to the Fermentas Catalog or go to www.fermentas.com/doubledigest to choose the best buffer for your experiments.

1X Tango™ Buffer:

33 mM Tris-acetate (pH 7.9 at 37°C), 10 mM magnesium acetate, 66 mM potassium acetate, 0.1 mg/ml BSA.

Storage Buffer

Eco52I is supplied in: 10 mM Tris-HCl (pH 8.2 at 25°C), 500 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 0.2 mg/ml BSA and 50% glycerol.

Recommended Protocol for Digestion

- Add:
nuclease-free water 16 µl
10X Buffer Eco52I 2 µl
DNA (0.5-1 µg/µl) 1 µl
Eco52I 0.5-2 µl
- Mix gently and spin down for a few seconds.
- Incubate at 37°C for 1-16 hours.

The digestion reaction may be scaled either up or down.

Recommended Protocol for Digestion of PCR Products Directly after Amplification

- Add:
PCR reaction mixture 10 µl (~0.1-0.5 µg of DNA)
nuclease-free water 18 µl
10X Buffer Eco52I 2 µl
Eco52I 1-2 µl
- Mix gently and spin down for a few seconds.
- Incubate at 37°C for 1-16 hours.

Thermal Inactivation

Eco52I is inactivated by incubation at 65°C for 20 min.

ENZYME PROPERTIES

Enzyme Activity in Fermentas REase Buffers, %

Eco52I	B	G	O	R	Tango™	2X Tango™
100	0-20	0-20	0-20	20-50	0-20	20-50

Methylation Effects on Digestion

Dam: never overlaps – no effect.
Dcm: never overlaps – no effect.
CpG: completely overlaps – blocked.
EcoKI: never overlaps – no effect.
EcoBI: never overlaps – no effect.

Stability during Prolonged Incubation

A minimum 0.2 units of the enzyme is required for complete digestion of 1 µg of lambda DNA in 16 hours at 37°C.

Digestion of Agarose-embedded DNA

A minimum of 5 units of the enzyme is required for complete digestion of 1 µg of agarose-embedded lambda DNA in 16 hours.

Compatible Ends

Bsp120I, CfrI, NotI

Number of Recognition Sites in DNA

λ	ΦX174	pBR322	pUC57	pUC18/19	pTZ19R/U	M13mp18/19
2	0	1	0	0	0	0

For **QUALITY CONTROL ASSAY DATA** see back page

QUALITY CONTROL ASSAY DATA

Overdigestion Assay

No detectable change in the specific fragmentation pattern is observed after a 320-fold overdigestion with Eco52I (20 u/μg lambda DNA x 16 hours).

Ligation/Recutting Assay

After a 50-fold overdigestion (3 u/μg DNA x 17 hours) with Eco52I, more than 95% of the digested DNA fragments can be ligated at a 5'-termini concentration of 0.01 μM. More than 95% of these sites can be recut.

Labeled Oligonucleotide (LO) Assay

No detectable degradation of single-stranded double-stranded labeled oligonucleotides occurred during incubation with 10 units of Eco52I for 4 hours.

Blue/White Cloning Assay

A mixture of pUC57/HindIII, pUC57/Eco32I and pUC57/PstI digests was incubated with 10 units of Eco52I for 16 hours. After religation and transformation, the background level of white colonies was 0.4%.

Quality authorized by:



Jurgita Zilinskiene

PRODUCT USE LIMITATION.

This product is developed, designed and sold exclusively *for research purposes and in vitro use only*. The product was not tested for use in diagnostics or for drug development, nor is it suitable for administration to humans or animals.

Please refer to www.fermentas.com for Material Safety Data Sheet of the product.