

CERTIFICATE OF ANALYSIS

Hinfl

#ER0802 5x2000 u

Lot: Expiry Date:

5'...G↓A N T C...3'
3'...C T N A↑G...5'

Concentration: 10 u/μl
Source: *Haemophilus influenzae* Rf
Supplied with: 2x1 ml of 10X Buffer R
 1 ml of 10X Buffer Tango™

Store at -20°C



In total 8 vials.

BSA included: Lot# BSA62-313P

RECOMMENDATIONS

1X Buffer R (for 100% Hinfl digestion)

10 mM Tris-HCl (pH 8.5), 10 mM MgCl₂, 100 mM KCl,
0.1 mg/ml BSA.

Incubation temperature

37°C.

Unit Definition

One unit is defined as the amount of Hinfl required to digest 1 μg lambda DNA in 1 hour at 37°C in 50 μl of recommended reaction buffer.

Dilution

Dilute with Dilution Buffer (#B19): 10 mM Tris-HCl (pH 7.4 at 25°C), 100 mM KCl, 1 mM EDTA, 1 mM DTT, 0.2 mg/ml BSA and 50% glycerol.

Double Digests

Tango™ Buffer is provided to simplify buffer selection for double digests. 98% of Fermentas Restriction Enzymes are active in a 1X or 2X concentration of Tango™ Buffer. Please refer to the Fermentas Catalog or go to www.fermentas.com/doubledigest to choose the best buffer for your experiments.

1X Tango™ Buffer:

33 mM Tris-acetate (pH 7.9), 10 mM magnesium acetate, 66 mM potassium acetate, 0.1 mg/ml BSA.

Storage Buffer

Hinfl is supplied in: 10 mM Tris-HCl (pH 7.5 at 25°C), 100 mM KCl, 1 mM DTT, 0.1 mM EDTA, 0.2 mg/ml BSA and 50% glycerol.

Recommended Protocol for Digestion

- Add:

nuclease-free water	16 µl
10X Buffer R	2 µl
DNA (0.5-1 µg/µl)	1 µl
Hinfl	0.5-2 µl*
- Mix gently and spin down for a few seconds.
- Incubate at 37°C for 1-16 hours.

The digestion reaction may be scaled either up or down.

Recommended Protocol for Digestion of PCR Products Directly after Amplification

- Add:

PCR reaction mixture	10 µl (~0.1-0.5 µg of DNA)
nuclease-free water	18 µl
10X Buffer R	2 µl
Hinfl	1-2 µl*
- Mix gently and spin down for a few seconds.
- Incubate at 37°C for 1-16 hours.

* This volume of the enzyme is recommended for preparations of standard concentrations (10 u/µl), whereas HC enzymes (50 u/µl) should be diluted with the Dilution Buffer to obtain 10 u/µl concentration.

Thermal Inactivation

Hinfl is inactivated by incubation at 65°C for 20 min.

ENZYME PROPERTIES

Enzyme Activity in Fermentas REase Buffers, %

B	G	O	R	Tango™	2X Tango™
0-20	20-50	50-100	100	50-100	50-100

Methylation Effects on Digestion

Dam: never overlaps – no effect.
Dcm: never overlaps – no effect.
CpG: may overlap – cleavage impaired.
EcoKI: never overlaps – no effect.
EcoBI: may overlap – blocked.

Stability during Prolonged Incubation

A minimum of 0.1 units of the enzyme is required for complete digestion of 1 µg of lambda DNA in 16 hours at 37°C.

Compatible Ends

G↓A(A/T)TC – Pfl.

Number of Recognition Sites in DNA

λ	ΦX174	pBR322	pUC57	pUC18/19	pTZ19R/U	M13mp18/19
148	21	10	5	6	9	27

For **QUALITY CONTROL ASSAY DATA** see back page

QUALITY CONTROL ASSAY DATA

Overdigestion Assay

No detectable change in the specific fragmentation pattern is observed after a 320-fold overdigestion with HinfI (20 u/μg lambda DNA x 16 hours).

Ligation/Recutting Assay

After a 50-fold overdigestion (3 u/μg DNA x 17 hours) with HinfI, more than 95% of the digested DNA fragments can be ligated at a 5'-termini concentration of 1.3 μM. More than 95% of these sites can be recut.

Labeled Oligonucleotide (LO) Assay

No detectable degradation of single-stranded or double-stranded labeled oligonucleotides occurred during incubation with 10 units of HinfI for 4 hours.

Quality authorized by:

 Jurgita Zilinskiene

PRODUCT USE LIMITATION.

This product is developed, designed and sold exclusively *for research purposes and in vitro use only*. The product was not tested for use in diagnostics or for drug development, nor is it suitable for administration to humans or animals.

Please refer to www.fermentas.com for Material Safety Data Sheet of the product.