

CERTIFICATE OF ANALYSIS

ApaI

#ER1412 5 x 3000 u

Lot: Expiry Date:

5'... **G G G C C**↓**C**...3'
3'... **C**↑**C C G G G**...5'

Concentration: 10 u/μl
Source: *E.coli* that carries the cloned *apaI*
 gene from *Acetobacter pasteurianus*
Supplied with: 2 x 1 ml of 10X Buffer B
 1 ml of 10X Buffer Tango™

Store at -20°C



In total 8 vials.

BSA included: Lot# BSA62-313P

RECOMMENDATIONS

1X Buffer B (for 100% ApaI digestion)

10 mM Tris-HCl (pH 7.5), 10 mM MgCl₂, 0.1 mg/ml BSA.

Incubation temperature

37°C*.

Unit Definition

One unit is defined as the amount of ApaI required to digest 1 μg of lambda DNA-Cpol fragments in 1 hour at 37°C in 50 μl of recommended reaction buffer.

Dilution

Dilute with Dilution Buffer (#B19): 10 mM Tris-HCl (pH7.4 at 25°C), 100 mM KCl, 1 mM EDTA, 1 mM DTT, 0.2 mg/ml BSA and 50% glycerol.

Double Digests

Tango™ Buffer is provided to simplify buffer selection for double digests. 98% of Fermentas restriction enzymes are active in a 1X or 2X concentration of Tango™ Buffer. Please refer to the Fermentas Catalog or go to www.fermentas.com/doubledigest to choose the best buffer for your experiments.

1X Tango™ Buffer:

33 mM Tris-acetate (pH 7.9), 10 mM magnesium acetate, 66 mM potassium acetate, 0.1 mg/ml BSA.

* Incubation at 30°C results in a 2-fold increase in activity.

Storage Buffer

Apal is supplied in: 10 mM Tris-HCl (pH 7.4 at 25°C), 50 mM NaCl, 1 mM DTT, 1 mM EDTA, 0.2 mg/ml BSA and 50% glycerol.

Recommended Protocol for Digestion

- Add:

nuclease-free water	16 µl
10X Buffer B	2 µl
DNA (0.5-1 µg/µl)	1 µl
Apal	0.5-2 µl
- Mix gently and spin down for a few seconds.
- Incubate at 37°C for 1-16 hours.

The digestion reaction may be scaled either up or down.

Recommended Protocol for Digestion of PCR Products Directly after Amplification

- Add:

PCR reaction mixture	10 µl (~0.1-0.5 µg of DNA)
nuclease-free water	18 µl
10X Buffer B	2 µl
Apal	1-2 µl
- Mix gently and spin down for a few seconds.
- Incubate at 37°C for 1-16 hours.

Thermal Inactivation

Apal is inactivated by incubation at 65°C for 20 min.

ENZYME PROPERTIES

Enzyme Activity in Fermentas REase Buffers, %

B	G	O	R	Tango™	2X Tango™
100	20-50	0-20	0-20	20-50	0-20

Methylation Effects on Digestion

Dam: never overlaps – no effect.
Dcm: may overlap – cleavage impaired.
CpG: may overlap – cleavage impaired.
EcoKI: never overlaps – no effect.
EcoBI: never overlaps – no effect.

Stability during Prolonged Incubation

A minimum of 0.2 units of the enzyme is required for complete digestion of 1 µg of lambda DNA in 16 hours at 37°C.

Digestion of Agarose-embedded DNA

A minimum of 5 units of the enzyme is required for complete digestion of 1 µg of agarose-embedded lambda DNA in 16 hours.

Compatible Ends

BseSI (GGGCC↓C), Eco24I (GGGCC↓C),
SduI (GGGCC↓C)

Number of Recognition Sites in DNA

λ	ΦX174	pBR322	pUC57	pUC18/19	pTZ19R/U	M13mp18/19
1	0	0	1	0	0	0

QUALITY CONTROL ASSAY DATA

Overdigestion Assay

No detectable change in the specific fragmentation pattern is observed after a 320-fold overdigestion with Apal (20 u/μg lambda DNA x 16 hours).

Ligation/Recutting Assay

After a 50-fold overdigestion (3 u/μg DNA x 17 hours) with Apal, more than 95% of the digested DNA fragments can be ligated at a 5'-termini concentration of 0.02 μM. More than 95% of these sites can be recut.

Labeled Oligonucleotide (LO) Assay

No detectable degradation of single-stranded or double-stranded labeled oligonucleotides occurred during incubation with 10 units of Apal for 4 hours.

Blue/White Cloning Assay

pUC57 was incubated with 10 units of Apal for 16 hours. After religation and transformation, the background level of white colonies was 0.1%.

Quality authorized by:



Jurgita Zilinskiene

This product is licensed under one or more U.S. Patents Nos. 5,179,015, 5,200,333, 5,320,957 or corresponding foreign patents.

Certain countries are out of the scope of patent coverage.

PRODUCT USE LIMITATION.

This product is developed, designed and sold exclusively *for research purposes and in vitro use only*. The product was not tested for use in diagnostics or for drug development, nor is it suitable for administration to humans or animals.

Please refer to www.fermentas.com for Material Safety Data Sheet of the product.